

2006

# On state-dependent delay partial neutral functional–differential equations

Eduardo Hernandez M.  
*ICMC, Universidade de Sao Paulo*

Mark A. McKibben  
*West Chester University of Pennsylvania, mmckibben@wcupa.edu*

Follow this and additional works at: [http://digitalcommons.wcupa.edu/math\\_facpub](http://digitalcommons.wcupa.edu/math_facpub)



Part of the [Partial Differential Equations Commons](#)

---

## Recommended Citation

Hernandez M., E., & McKibben, M. A. (2006). On state-dependent delay partial neutral functional–differential equations. *Applied Mathematics and Computation* Retrieved from [http://digitalcommons.wcupa.edu/math\\_facpub/4](http://digitalcommons.wcupa.edu/math_facpub/4)

This Article is brought to you for free and open access by the College of Arts & Sciences at Digital Commons @ West Chester University. It has been accepted for inclusion in Mathematics by an authorized administrator of Digital Commons @ West Chester University. For more information, please contact [wcressler@wcupa.edu](mailto:wcressler@wcupa.edu).



# On state-dependent delay partial neutral functional–differential equations

Eduardo Hernández M.<sup>a,\*</sup>, Mark A. McKibben<sup>b</sup>

<sup>a</sup> *Departamento de Matemática, ICMC, Universidade de São Paulo, Caixa Postal 668, 13560-970 São Carlos SP, Brazil*

<sup>b</sup> *Department of Mathematics and Computer Science, Goucher College, 1021 Dulaney Valley Road, Baltimore, MD 21204, USA*

## Abstract

In this paper, we study the existence of mild solutions for a class of abstract partial neutral functional–differential equations with state-dependent delay.

© 2006 Elsevier Inc. All rights reserved.

*Keywords:* Abstract Cauchy problem; Neutral equations; State-dependent delay; Semigroup of linear operators; Unbounded delay

## 1. Introduction

The purpose of this article is establish the existence of mild solutions for a class of abstract neutral functional–differential equations with state-dependent delay described by the form

$$\frac{d}{dt}D(u_t) = AD(u_t) + F(t, x_{\rho(t, x_t)}), \quad t \in I = [0, a], \quad (1.1)$$

$$x_0 = \varphi \in \mathcal{B}, \quad (1.2)$$

where  $A$  is the infinitesimal generator of a compact  $C_0$ -semigroup of bounded linear operators  $(T(t))_{t \geq 0}$  on a Banach space  $X$ ; the function  $x_s : (-\infty, 0] \rightarrow X$ ,  $x_s(\theta) = x(s + \theta)$ , belongs to some abstract phase space  $\mathcal{B}$  described axiomatically;  $F$ ,  $G$  are appropriate functions; and  $D\psi = \psi(0) - G(t, \psi)$ , where  $\psi$  is in  $\mathcal{B}$ .

Functional–differential equations with state-dependent delay appear frequently in applications as model of equations and for this reason the study of this type of equation has received a significant amount of attention in the last years, see for instance [1–11] and the references therein. We also cite [12,9,13] for the case of neutral differential equations with dependent delay. The literature related to partial functional–differential equations with state-dependent delay is limited, to our knowledge, to the recent works [14,15].

Abstract neutral differential equations arise in many areas of applied mathematics. For this reason, they have largely been studied during the last few decades. The literature related to ordinary neutral differential

\* Corresponding author.

*E-mail addresses:* [lalohm@icmc.usp.br](mailto:lalohm@icmc.usp.br) (E. Hernández M.), [mmckibben@goucher.edu](mailto:mmckibben@goucher.edu) (M.A. McKibben).

equations is very extensive, thus, we refer the reader to [16] only, which contains a comprehensive description of such equations. Similarly, for more on partial neutral functional–differential equations and related issues we refer to Adimy and Ezzinbi [17], Hale [18], Wu and Xia [19] and [20] for finite delay equations, and Hernández and Henriquez [21,22] and Hernández [23] for unbounded delays.

## 2. Preliminaries

Throughout this paper,  $A: D(A) \subset X \rightarrow X$  is the infinitesimal generator of a compact  $C_0$ -semigroup of linear operators  $(T(t))_{t \geq 0}$  on a Banach space  $X$  and  $\tilde{M}$  is a positive constant such that  $\|T(t)\| \leq \tilde{M}$  for every  $t \in I = [0, a]$ . For background information related to semigroup theory, we refer the reader to Pazy [24].

In this work we will employ an axiomatic definition for the phase space  $\mathcal{B}$  which is similar to those introduced in [25]. Specifically,  $\mathcal{B}$  will be a linear space of functions mapping  $(-\infty, 0]$  into  $X$  endowed with a seminorm  $\|\cdot\|_{\mathcal{B}}$ , and satisfies the following axioms:

- (A) If  $x: (-\infty, \sigma + b] \rightarrow X$ ,  $b > 0$ , is such that  $x|_{[\sigma, \sigma + b]} \in C([\sigma, \sigma + b]: X)$  and  $x_\sigma \in \mathcal{B}$ , then for every  $t \in [\sigma, \sigma + b]$  the following conditions hold:
- (i)  $x_t$  is in  $\mathcal{B}$ ,
  - (ii)  $\|x(t)\| \leq H\|x_t\|_{\mathcal{B}}$ ,
  - (iii)  $\|x_t\|_{\mathcal{B}} \leq K(t - \sigma) \sup\{\|x(s)\| : \sigma \leq s \leq t\} + M(t - \sigma)\|x_\sigma\|_{\mathcal{B}}$ ,  
where  $H > 0$  is a constant;  $K, M: [0, \infty) \rightarrow [1, \infty)$ ,  $K$  is continuous,  $M$  is locally bounded, and  $H, K, M$  are independent of  $x(\cdot)$ .
- (A1) For the function  $x(\cdot)$  in (A), the function  $t \rightarrow x_t$  is continuous from  $[\sigma, \sigma + b]$  into  $\mathcal{B}$ .
- (B) The space  $\mathcal{B}$  is complete.

**Example 2.1** (The phase spaces  $C_g, C_g^0$ ). Let  $g: (-\infty, 0] \rightarrow [1, \infty)$  be a continuous, non-decreasing function with  $g(0) = 1$ , which satisfies conditions (g-1), (g-2) of [25]. Briefly, this means that the function  $\gamma(t) := \sup_{-\infty < \theta \leq -t} \frac{g(t+\theta)}{g(\theta)}$  is locally bounded for  $t \geq 0$  and that  $g(\theta) \rightarrow \infty$  as  $\theta \rightarrow -\infty$ .

Let  $C_g(X)$  be the vector space consisting of the continuous functions  $\varphi$  such that  $\frac{\varphi}{g}$  is bounded on  $(-\infty, 0]$ , and let  $C_g^0(X)$  be the subspace of  $C_g(X)$  containing precisely those functions  $\varphi$  for which  $\frac{\varphi(\theta)}{g(\theta)} \rightarrow 0$  as  $\theta \rightarrow -\infty$ . The spaces  $C_g$  and  $C_g^0$ , endowed with the norm  $\|\varphi\|_g := \sup_{\theta \leq 0} \frac{\|\varphi(\theta)\|}{g(\theta)}$ , are both phase spaces which satisfy axioms (A), (A-1), (B), see [25, Theorem 1.3.6] for details. Moreover, in this case  $K(t) = 1$ , for every  $t \geq 0$ .

**Example 2.2** (The phase space  $C_r \times L^p(g; X)$ ). Assume that  $g: (-\infty, -r) \rightarrow \mathbb{R}$  is a Lebesgue integrable function and that there exists a non-negative and locally bounded function  $\gamma$  such that  $g(\xi + \theta) \leq \gamma(\xi) g(\theta)$ , for all  $\xi \leq 0$  and  $\theta \in (-\infty, -r) \setminus N_\xi$ , where  $N_\xi \subseteq (-\infty, -r)$  is a set with Lebesgue measure zero. The space  $C_r \times L^p(g; X)$  consists of all classes of functions  $\varphi: (-\infty, 0] \rightarrow X$  such that  $\varphi$  is continuous on  $[-r, 0]$  and  $g^{\frac{1}{p}}\|\varphi\| \in L^p((-\infty, -r); X)$ . The seminorm in  $C_r \times L^p(g; X)$  is defined by

$$\|\varphi\|_{\mathcal{B}} := \sup\{\|\varphi(\theta)\| : -r \leq \theta \leq 0\} + \left( \int_{-\infty}^{-r} g(\theta) \|\varphi(\theta)\|^p d\theta \right)^{1/p}.$$

If  $g(\cdot)$  satisfies the conditions (g-5), (g-6) and (g-7) in the nomenclature of [25], then  $\mathcal{B} = C_r \times L^p(g; X)$  satisfies axioms (A), (A1), (B) (see [25, Theorem 1.3.8] for details). Moreover, when  $r = 0$  and  $p = 2$ , we have that  $H = 1$ ,  $M(t) = \gamma(-t)^{1/2}$ , and  $K(t) = 1 + (\int_{-t}^0 g(\theta) d\theta)^{1/2}$  for  $t \geq 0$ .

**Remark 2.1.** Let  $\varphi \in \mathcal{B}$  and  $t \leq 0$ . The notation  $\varphi_t$  represents the function  $\varphi_t: (-\infty, 0] \rightarrow X$  defined by  $\varphi_t(\theta) = \varphi(t + \theta)$ . Consequently, if the function  $x(\cdot)$  in axiom (A) is such that  $x_0 = \varphi$ , then  $x_t = \varphi_t$ . We observe that  $\varphi_t$  is well defined for every  $t < 0$  since the domain of  $\varphi(\cdot)$  is  $(-\infty, 0]$ .

We also note that, in general,  $\varphi_t \notin \mathcal{B}$ ; consider, for example, the characteristic function  $\mathcal{X}_{[\mu, 0]}$ ,  $\mu < -r < 0$ , in the space  $C_r \times L^p(g; X)$ .

The terminology and notations are those generally used in functional analysis. In particular, for Banach spaces  $(Z, \|\cdot\|_Z)$ ,  $(W, \|\cdot\|_W)$ , the notation  $\mathcal{L}(Z; W)$  stands for the Banach space of bounded linear operators from  $Z$  into  $W$ , and we abbreviate this notation to  $\mathcal{L}(Z)$  when  $Z = W$ . Moreover,  $B_r(x, Z)$  denotes the closed ball with center at  $x$  and radius  $r > 0$  in  $Z$  and for a bounded function  $\xi: I \rightarrow Z$  and  $t \in [0, a]$  we employ the notation  $\|\xi\|_{Z,t}$  for

$$\|\xi\|_{Z,t} = \sup\{\|\xi(s)\|_Z : s \in [0, t]\}. \quad (2.3)$$

We will simply write  $\|\xi\|_t$  when no confusion arises.

The remainder of the paper is divided into two sections. The existence of mild solutions for the abstract Cauchy problem (1.1)–(1.2) is studied in Section 3, and Section 4 is devoted to a discussion of some applications. To conclude the current section, we recall the following well-known result, referred to as the Leray Schauder Alternative, for convenience.

**Theorem 2.1.** [26, Theorem 6.5.4] *Let  $D$  be a convex subset of a Banach space  $X$  and assume that  $0 \in D$ . Let  $G: D \rightarrow D$  be a completely continuous map. Then, either the set  $\{x \in D: x = \lambda G(x), 0 < \lambda < 1\}$  is unbounded or the map  $G$  has a fixed point in  $D$ .*

### 3. Existence results

In this section we discuss the existence of mild solutions for the abstract system (1.1)–(1.2). We begin by introducing the following conditions:

(H <sub>$\varphi$</sub> ) Let  $\mathcal{R}(\rho^-) = \{\rho(s, \psi) : (s, \psi) \in I \times \mathcal{B}, \rho(s, \psi) \leq 0\}$ . The function  $t \rightarrow \varphi_t$  is well defined and continuous from  $\mathcal{R}(\rho^-)$  into  $\mathcal{B}$ , and there exists a continuous and bounded function  $J^\varphi: \mathcal{R}(\rho^-) \rightarrow (0, \infty)$  such that  $\|\varphi_t\|_{\mathcal{B}} \leq J^\varphi(t)\|\varphi\|_{\mathcal{B}}$  for every  $t \in \mathcal{R}(\rho^-)$ .

(H<sub>1</sub>) The function  $F: I \times \mathcal{B} \rightarrow X$  satisfies the following properties:

- The function  $F(\cdot, \psi): I \rightarrow X$  is strongly measurable, for every  $\psi \in \mathcal{B}$ .
- The function  $F(t, \cdot): \mathcal{B} \rightarrow X$  is continuous, for each  $t \in I$ .
- There exist a continuous non decreasing function  $W: [0, \infty) \rightarrow (0, \infty)$  and an integrable function  $m: I \rightarrow [0, \infty)$  such that

$$\|F(t, \psi)\| \leq m(t)W(\|\psi\|_{\mathcal{B}}), \quad (t, \psi) \in I \times \mathcal{B}.$$

(H<sub>2</sub>) The function  $G: \mathbb{R} \times \mathcal{B} \rightarrow X$  is continuous and there exists  $L_G > 0$  such that

$$\|G(t, \psi_1) - G(t, \psi_2)\| \leq L_G\|\psi_1 - \psi_2\|_{\mathcal{B}}, \quad (t, \psi_i) \in I \times \mathcal{B}.$$

(H<sub>3</sub>) Let  $S(a) = \{x: (-\infty, a] \rightarrow X: x_0 = 0; x \in C([0, a]: X)\}$  endowed with the norm of uniform convergence on  $[0, a]$  and  $y: (-\infty, a] \rightarrow X$  be the function defined by  $y_0 = \varphi$  on  $(-\infty, 0]$  and  $y(t) = T(t)\varphi(0)$  on  $[0, a]$ . Then, for every bounded set  $Q$  such that  $Q \subset S(a)$ , the set of functions  $\{t \rightarrow G(t, x_t + y_t): x \in Q\}$  is equicontinuous on  $[0, a]$ .

**Remark 3.2.** We remark that condition H <sub>$\varphi$</sub>  is frequently satisfied by functions that are continuous and bounded. In fact, if the space  $\mathcal{B}$  satisfies axiom C<sub>2</sub> in [25], then there exists a constant  $L > 0$  such that  $\|\varphi_t\|_{\mathcal{B}} \leq L \sup_{\theta \leq 0} \|\varphi(\theta)\|$  for every  $\varphi \in \mathcal{B}$  that is continuous and bounded, see [25, Proposition 7.1.1] for details. Consequently,

$$\|\varphi_t\|_{\mathcal{B}} \leq L \frac{\sup_{\theta \leq 0} \|\varphi(\theta)\|}{\|\varphi\|_{\mathcal{B}}} \|\varphi\|_{\mathcal{B}}$$

for every  $\varphi \in \mathcal{B} \setminus \{0\}$  continuous and bounded and every  $t \leq 0$ . We also observe that  $C_r \times L^p(g; X)$  satisfies axiom C<sub>2</sub> if  $g(\cdot)$  is integrable on  $(-\infty, r]$ , see [25, p. 10].

Motivated by general semigroup theory, we adopt the following concept of mild solution.

**Definition 3.1.** A function  $x: (-\infty, a] \rightarrow X$  is a mild solution of the abstract Cauchy problem (1.1)–(1.2) if  $x_0 = \varphi$ ,  $x_{\rho(s, x_s)} \in \mathcal{B}$ , for every  $s \in I$ , and

$$x(t) = T(t)(\varphi(0) - G(0, \varphi)) + G(t, x_t) + \int_0^t T(t-s)F(s, x_{\rho(s, x_s)}) ds, \quad t \in I.$$

The proof of Lemma 3.1 is routine; the details are left to the reader.

**Lemma 3.1.** Let  $x: (-\infty, a] \rightarrow X$  be a function such that  $x_0 = \varphi$  and  $x|_{[0, a]} \in C([0, a] : X)$ . Then

$$\|x_s\|_{\mathcal{B}} \leq (M_a + J_0^\varphi) \|\varphi\|_{\mathcal{B}} + K_a \sup\{\|x(\theta)\|; \theta \in [0, \max\{0, s\}]\}, \quad s \in \mathcal{R}(\rho^-) \cup [0, a],$$

where  $J_0^\varphi = \sup_{t \in \mathcal{R}(\rho^-)} J^\varphi(t)$  and the notation in (2.3) has been used.

Now, we can prove our first existence result.

**Theorem 3.2.** Let  $\varphi \in \mathcal{B}$  and assume that conditions  $H_1$ ,  $H_2$ ,  $H_\varphi$  hold. If

$$K_a \left( L_G + \tilde{M} \liminf_{\xi \rightarrow \infty^+} \frac{W(\xi)}{\xi} \int_0^a m(s) ds \right) < 1,$$

then there exists a mild solution of (1.1)–(1.2).

**Proof.** Consider the space  $Y = \{u \in C(I; X) : u(0) = \varphi(0)\}$  endowed with the uniform convergence topology, and define the operator  $\Gamma : Y \rightarrow Y$  by

$$\Gamma x(t) = T(t)(\varphi(0) - G(0, \varphi)) + G(t, \bar{x}_t) + \int_0^t T(t-s)F(s, \bar{x}_{\rho(s, \bar{x}_s)}) ds, \quad t \in I,$$

where  $\bar{x} : (-\infty, a] \rightarrow X$  is the extension of  $x$  to  $(-\infty, a]$  such that  $\bar{x}_0 = \varphi$ . From our assumptions it is easy to see that  $\Gamma x \in Y$ .

Let  $\bar{\varphi} : (-\infty, a] \rightarrow X$  be the extension of  $\varphi$  to  $(-\infty, a]$  such that  $\bar{\varphi}(\theta) = \varphi(0)$  on  $I$ . We claim that there exists  $r > 0$  such that  $\Gamma(B_r(\bar{\varphi}|_I, Y)) \subset B_r(\bar{\varphi}|_I, Y)$ . Indeed, suppose to the contrary that this assertion is false. Then, for every  $r > 0$  there exist  $x^r \in B_r(\bar{\varphi}|_I, Y)$  and  $t^r \in I$  such that  $r < \|\Gamma x^r(t^r) - \varphi(0)\|$ . Then, from Lemma 3.1 we find that

$$\begin{aligned} r &< \|\Gamma x^r(t^r) - \varphi(0)\| \\ &\leq \|T(t^r)\varphi(0) - \varphi(0)\| + \|T(t^r)G(0, \varphi) - G(0, \varphi)\| + \|G(t^r, \overline{(x^r)_{t^r}}) - G(0, \varphi)\| \\ &\quad + \int_0^{t^r} \|T(t^r-s)\| \|F(s, \overline{(x^r)_{\rho(s, \overline{(x^r)_{t^r}})}})\| ds \\ &\leq (\tilde{M} + 1)H \|\varphi\|_{\mathcal{B}} + \|T(t^r)G(0, \varphi) - G(0, \varphi)\| + L_G(K_a \|x^r - \varphi(0)\|_{t^r} + (M_a + HK_a + 1)\|\varphi\|) \\ &\quad + \tilde{M} \int_0^{t^r} m(s)W((M_a + J_0^\varphi)\|\varphi\|_{\mathcal{B}} + K_a(\|x^r - \varphi(0)\|_a + \|\varphi(0)\|)) ds \\ &\leq ((\tilde{M} + 1)H)\|\varphi\|_{\mathcal{B}} + \|T(t^r)G(0, \varphi) - G(0, \varphi)\| + L_G(K_a r + (M_a + HK_a + 1)\|\varphi\|) \\ &\quad + \tilde{M} \int_0^{t^r} m(s)W((M_a + J_0^\varphi + H)\|\varphi\|_{\mathcal{B}} + K_a r) ds \end{aligned}$$

and hence

$$1 \leq K_a \left( L_G + \tilde{M} \liminf_{\xi \rightarrow \infty} \frac{W(\xi)}{\xi} \int_0^a m(s) ds \right),$$

which contradicts our assumption.

Let  $r > 0$  be such that  $\Gamma(B_r(\bar{\varphi}|_I, Y)) \subset B_r(\bar{\varphi}|_I, Y)$  and consider the decomposition  $\Gamma = \Gamma_1 + \Gamma_2$  where

$$\Gamma_1 x(t) = T(t)(\varphi(0) - G(0, \varphi)) + G(t, \bar{x}_t), \quad t \in I,$$

$$\Gamma_2 x(t) = \int_0^t T(t-s)F(s, \bar{x}_{\rho(s, \bar{x}_s)}) ds, \quad t \in I.$$

From the proof of [14, Theorem 2.2], we know that  $\Gamma_2$  is completely continuous. Moreover, using the phase space axioms we find that

$$\|\Gamma_1 u(t) - \Gamma_1 v(t)\| \leq L_G K_a \|u - v\|_a, \quad t \in I,$$

which proves that  $\Gamma_1$  is a contraction on  $B_r(\bar{\varphi}|_I, Y)$ , so that  $\Gamma$  is a condensing operator on  $B_r(\bar{\varphi}|_I, Y)$ .

The existence of a mild solution for (1.1)–(1.2) is now a consequence of [27, Theorem 4.3.2]. This completes the proof.  $\square$

**Theorem 3.3.** *Let conditions  $H_\varphi, H_1, H_3$  be satisfied. Assume that  $\rho(t, \psi) \leq t$  for every  $(t, \psi) \in I \times \mathcal{B}$ , that  $G$  is completely continuous, and that there exist positive constants  $c_1, c_2$  such that  $\|G(t, \psi)\| \leq c_1 \|\psi\|_{\mathcal{B}} + c_2$  for every  $(t, \psi) \in I \times \mathcal{B}$ . If  $\mu = 1 - K_a c_1 > 0$  and*

$$\frac{\tilde{M}K_a}{\mu} \int_0^a m(s) ds < \int_C \frac{ds}{W(s)},$$

where

$$C = \left( M_a + J_0^\varphi + \tilde{M}HK_a \right) \|\varphi\| + \frac{K_a}{\mu} \left[ \tilde{M}\|G(0, \varphi)\| + c_1(M_a + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + c_2 \right],$$

then there exists a mild solution of (1.1)–(1.2).

**Proof.** On the space  $\mathcal{BC} = \{u : (-\infty, a] \rightarrow X; u_0 = 0, u|_I \in C(I; X)\}$  endowed with the norm  $\|u\|_a = \sup_{s \in [0, a]} \|u(s)\|$ , we define the operator  $\Gamma : \mathcal{BC} \rightarrow \mathcal{BC}$  by

$$\Gamma x(t) = \begin{cases} 0, & t \in (-\infty, 0], \\ T(t)G(0, \varphi) - G(t, \bar{x}_t) + \int_0^t T(t-s)F(s, \bar{x}_{\rho(s, \bar{x}_s)}) ds, & t \in I, \end{cases}$$

where  $\bar{x} : (-\infty, a] \rightarrow X$  is defined by the relation  $\bar{x} = y + x$  on  $(-\infty, a]$ . In preparation for using Theorem 2.1, we establish *a priori* estimates for the solutions of the integral equation  $z = \lambda \Gamma z$ ,  $\lambda \in (0, 1)$ . Let  $x^\lambda$  be a solution of  $z = \lambda \Gamma z$ ,  $\lambda \in (0, 1)$ . If  $\alpha^\lambda(s) = \sup_{\theta \in [0, s]} \|x^\lambda(\theta)\|$ , then from Lemma 3.1 and the fact that  $\rho(s, (\bar{x}^\lambda)_s) \leq s$ , we find that

$$\begin{aligned} \|x^\lambda(t)\| &\leq \|T(t)G(0, \varphi)\| + c_1 \|\bar{x}_t\|_{\mathcal{B}} + c_2 + \tilde{M} \int_0^t m(s)W((M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha^\lambda(s)) ds \\ &\leq \tilde{M}\|G(0, \varphi)\| + c_1((M_a + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha(t)) + c_2 \\ &\quad + \tilde{M} \int_0^t m(s)W((M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha^\lambda(s)) ds. \end{aligned}$$

Consequently,

$$\begin{aligned} \|\alpha^\lambda(t)\| &\leq \tilde{M}\|G(0, \varphi)\| + c_1((M_a + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha(t)) + c_2 \\ &\quad + \tilde{M} \int_0^t m(s)W((M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha^\lambda(s)) ds \end{aligned}$$

and so

$$\begin{aligned} \|\alpha^\lambda(t)\| &\leq \frac{1}{\mu} [\tilde{M}\|G(0, \varphi)\| + c_1(M_a + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + c_2] \\ &\quad + \frac{\tilde{M}}{\mu} \int_0^t m(s)W((M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha^\lambda(s)) ds. \end{aligned}$$

By denoting  $\xi^\lambda(t) = (M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + K_a \alpha(t)$ , we obtain after a rearrangement of terms that

$$\begin{aligned} \xi^\lambda(t) &\leq (M_a + J_0^\varphi + \tilde{M}HK_a)\|\varphi\| + \frac{K_a}{\mu} [\tilde{M}\|G(0, \varphi)\| + c_1(M_a + \tilde{M}HK_a)\|\varphi\|_{\mathcal{B}} + c_2] \\ &\quad + \frac{\tilde{M}K_a}{\mu} \int_0^t m(s)W(\xi^\lambda(s)) ds. \end{aligned} \tag{3.4}$$

Denoting by  $\beta_\lambda(t)$  the right-hand side of (3.4), it follows that

$$\beta'_\lambda(t) \leq \frac{\tilde{M}K_a}{\mu} m(t)W(\beta_\lambda(t))$$

and hence

$$\int_{\beta_\lambda(0)=c}^{\beta_\lambda(t)} \frac{ds}{W(s)} \leq \frac{\tilde{M}K_a}{\mu} \int_0^a m(s) ds < \int_c^\infty \frac{ds}{W(s)},$$

which implies that the set of functions  $\{\beta_\lambda(\cdot) : \lambda \in (0, 1)\}$  is bounded in  $C(I : \mathbb{R})$ . Thus, the set  $\{x^\lambda(\cdot) : \lambda \in (0, 1)\}$  is bounded on  $I$ .

To prove that  $\Gamma$  is completely continuous, we consider the decomposition  $\Gamma = \Gamma_1 + \Gamma_2$  introduced in the proof of Theorem 3.2. From the proof of [14, Theorem 2.2] we know that  $\Gamma_2$  is completely continuous and from the assumptions on  $G$  we infer that  $\Gamma_1$  is a compact map. It remains to show that  $\Gamma_1$  is continuous. Let  $(u^n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{BC}$  and  $u \in \mathcal{BC}$  such that  $u^n \rightarrow u$ . From the phase space axioms we infer that  $(\overline{u^n})_s \rightarrow \overline{u}_s$  uniformly on  $[0, a]$  as  $n \rightarrow \infty$  and that  $U = [0, a] \times \{(\overline{u^n})_s, \overline{u}_s : s \in [0, a], n \in \mathbb{N}\}$  is relatively compact in  $[0, a] \times \mathcal{B}$ . Thus,  $G$  is uniformly continuous on  $U$ , so that  $G(s, (\overline{u^n})_s) \rightarrow G(s, \overline{u}_s)$  uniformly on  $[0, a]$  as  $n \rightarrow \infty$ , which shows that  $\Gamma_1$  is continuous.

These remarks, in conjunction with Theorem 2.1, enable us to conclude that there exists a mild solution for (1.1) and (1.2). The proof is complete.  $\square$

#### 4. Examples

We conclude this work with two applications of our previous abstract results. In the sequel,  $X = L^2([0, \pi])$  and  $A : D(A) \subset X \rightarrow X$  is the operator  $Af = f''$  with domain  $D(A) := \{f \in X : f'' \in X, f(0) = f(\pi) = 0\}$ . It is well known that  $A$  is the infinitesimal generator of a compact  $C_0$ -semigroup of bounded linear operators  $(T(t))_{t \geq 0}$  on  $X$ . Moreover,  $A$  has discrete spectrum, the eigenvalues are  $-n^2$ ,  $n \in \mathbb{N}$ , with corresponding normalized eigenvectors  $z_n(\xi) := (\frac{2}{\pi})^{1/2} \sin(n\xi)$ , the set  $\{z_n : n \in \mathbb{N}\}$  is an orthonormal basis of  $X$ , and  $T(t)x = \sum_{n=1}^\infty e^{-n^2 t} \langle x, z_n \rangle z_n$  for  $x \in X$ . Consider the differential system

$$\begin{aligned} & \frac{d}{dt} \left[ u(t, \xi) + \int_{-\infty}^t a_1(s-t)u(s, \xi) ds \right] \\ &= \frac{\partial^2}{\partial \xi^2} \left[ u(t, \xi) + \int_{-\infty}^t a_1(s-t)u(s, \xi) ds \right] \\ & \quad + \int_{-\infty}^t a_2(s-t)u(s - \rho_1(t)\rho_2(\|u(t)\|), \xi) ds, \quad t \in I = [0, a], \quad \xi \in [0, \pi], \end{aligned} \quad (4.5)$$

together with the initial conditions

$$u(t, 0) = u(t, \pi) = 0, \quad t \geq 0, \quad (4.6)$$

$$u(\tau, \xi) = \varphi(\tau, \xi), \quad \tau \leq 0, \quad 0 \leq \xi \leq \pi. \quad (4.7)$$

In the sequel,  $\mathcal{B} = C_0 \times L^2(g; X)$  is the space introduced in Example 2.2;  $\varphi \in \mathcal{B}$  with the identification  $\varphi(s)(\tau) = \varphi(s, \tau)$ ; the functions  $a_i : \mathbb{R} \rightarrow \mathbb{R}$ ,  $\rho_i : [0, \infty) \rightarrow [0, \infty)$ ,  $i = 1, 2$ , are continuous; and

$$L_i = \left( \int_{-\infty}^0 \frac{(a_i(s))^2}{g(s)} ds \right)^{1/2} < 1, \quad i = 1, 2.$$

By defining the operators  $D, G, F : I \times \mathcal{B} \rightarrow X$  and  $\rho : I \times \mathcal{B} \rightarrow \mathbb{R}$  by

$$D(\psi) = \psi(0, \xi) - G(\psi)(\xi)$$

$$G(\psi)(\xi) = - \int_{-\infty}^0 a_1(s)\psi(s, \xi) ds,$$

$$F(t, \psi)(\xi) = \int_{-\infty}^0 a_2(s) \psi(s, \xi) ds,$$

$$\rho(s, \psi) = s - \rho_1(s) \rho_2(\|\psi(0)\|),$$

we can transform system (4.5)–(4.7) into the abstract system (1.1)–(1.2). Moreover,  $G, F$  are bounded linear operators,  $\|G\|_{\mathcal{L}(\mathcal{B}, X)} \leq L_1$  and  $\|F\|_{\mathcal{L}(\mathcal{B}, X)} \leq L_2$ .

The next result is an immediate consequence of Theorem 3.2.

**Theorem 4.4.** *Let  $\varphi \in \mathcal{B}$  be such that condition  $H_\varphi$  holds and assume that*

$$\left(1 + \left(\int_{-a}^0 g(\theta) d\theta\right)^{1/2}\right)(L_1 + aL_2) < 1. \quad (4.8)$$

Then there exists a mild solution of (4.5)–(4.7).

The proof of Corollary 4.1 follows directly from Theorem 4.4 and Remark 3.2.

**Corollary 4.1.** *Let  $\varphi \in \mathcal{B}$  continuous and bounded, and assume that (4.8) holds. Then, there exists a mild solution of (4.5)–(4.7) on  $I$ .*

To conclude this section, we briefly consider the differential system

$$\frac{d}{dt}[u(t, \xi) + u(t - r, \xi)] = \frac{\partial^2}{\partial \xi^2}[u(t, \xi) + u(t - r, \xi)] + a_1(t)b_1(u(t - \sigma(\|u(t)\|), \xi)),$$

$$t \in I = [0, a], \quad \xi \in [0, \pi], \quad (4.9)$$

$$u(t, 0) = u(t, \pi) = 0, \quad (4.10)$$

$$u(\tau, \xi) = \varphi(\tau, \xi), \quad \tau \leq 0, \quad \xi \in [0, \pi]. \quad (4.11)$$

For this system, we take  $\varphi \in \mathcal{B} = C_g^0(X)$  and assume that the functions  $a_1 : I \rightarrow \mathbb{R}$ ,  $b_1 : \mathbb{R} \times J \rightarrow \mathbb{R}$ ,  $\sigma : \mathbb{R} \rightarrow \mathbb{R}^+$  are continuous and that there exist positive constants  $d_1, d_2$  such that  $|b_1(t)| \leq d_1|t| + d_2$  for every  $t \in \mathbb{R}$ .

Let  $D, G : \mathcal{B} \rightarrow X$ ,  $F : [0, a] \times \mathcal{B} \rightarrow X$  and  $\rho : [0, a] \times \mathcal{B} \rightarrow \mathbb{R}$  be the operators defined by  $D(\psi)(\xi) = \psi(0, \xi) - G(\psi)(\xi)$ ,  $G(\psi)(\xi) = -\psi(-r, \xi)$ ,  $F(t, \psi)(\xi) = a_1(t)b_1(\psi(0, \xi))$  and  $\rho(t, \psi) = t - \sigma(\|\psi(0)\|)$ . Using these definitions, we can represent the system (4.8)–(4.10) in the abstract form (1.1) and (1.2). Moreover,  $G$  is a bounded linear operator on  $\mathcal{B}$  with  $\|G(\psi)\|_{\mathcal{L}(\mathcal{B}, X)} \leq g(-r)$ ;  $F$  is continuous and  $\|F(t, \psi)\| \leq a_1(t)[d_1\|\psi\|_{\mathcal{B}} + d_2\pi]$  for all  $(t, \psi) \in I \times \mathcal{B}$ . As such, the following results follow from Theorem 3.2 and Remark 3.2.

**Theorem 4.5.** *If  $\varphi \in \mathcal{B}$  satisfies condition  $H_\varphi$  and  $g(-r) + d_1 \int_0^a a_1(s) ds < 1$ , then there exists a mild solution of (4.9)–(4.11).*

**Corollary 4.2.** *If  $\varphi$  is continuous and bounded on  $(-\infty, 0]$  and  $g(-r) + d_1 \int_0^a a_1(s) ds < 1$ , then there exists a mild solution of (4.9)–(4.11).*

## References

- [1] Ovide Arino, Khalid Boushaba, Ahmed Boussouar, A mathematical model of the dynamics of the phytoplankton-nutrient system: spatial heterogeneity in ecological models (Alcalá de Henares, 1998), *Nonlinear Anal. RWA* 1 (1) (2000) 69–87.
- [2] Walter Aiello, H.I. Freedman, J. Wu, Analysis of a model representing stage-structured population growth with state-dependent time delay, *SIAM J. Appl. Math.* 52 (3) (1992) 855–869.
- [3] Maria Bartha, Periodic solutions for differential equations with state-dependent delay and positive feedback, *Nonlinear Anal. TMA* 53 (6) (2003) 839–857.
- [4] Yulin Cao, Jiangping Fan, Thomas C. Gard, The effects of state-dependent time delay on a stage-structured population growth model, *Nonlinear Anal. TMA* 19 (2) (1992) 95–105.
- [5] Domoshnitsky Alexander, Michael Drakhlin, Elena Litsyn, On equations with delay depending on solution, *Nonlinear Anal. TMA* 49 (5) (2002) 689–701.



- [6] Yongkun Li, Lifei Zhu, Positive periodic solutions for a class of higher-dimensional state-dependent delay functional differential equations with feedback control, *Appl. Math. Comput.* 159 (3) (2004) 783–795.
- [7] A.A.S. Zaghrout, S.H. Attalah, Analysis of a model of stage-structured population dynamics growth with time state-dependent time delay, *Appl. Math. Comput.* 77 (2–3) (1996) 185–194.
- [8] Ferenc Hartung, Parameter estimation by quasilinearization in functional differential equations with state-dependent delays: a numerical study, in: *Proceedings of the Third World Congress of Nonlinear Analysis, Part 7 (Catania, 2000)*, *Nonlinear Anal. TMA* 47 (7) (2001) 4557–4566.
- [9] Ferenc Hartung, Terry L. Herdman, Janos Turi, Parameter identification in classes of neutral differential equations with state-dependent delays, *Nonlinear Anal. TMA* 39 (3) (2000) 305–325.
- [10] Ferenc Hartung, Janos Turi, Identification of parameters in delay equations with state-dependent delays, *Nonlinear Anal. TMA* 29 (11) (1997) 1303–1318.
- [11] Y. Kuang, H.L. Smith, Slowly oscillating periodic solutions of autonomous state-dependent delay equations, *Nonlinear Anal. TMA* 19 (9) (1992) 855–872.
- [12] R.D. Driver, A neutral system with state-dependent delay, *J. Differen. Equat.* 54 (1) (1984) 73–86.
- [13] Zhihui Yang, Jinde Cao, Existence of periodic solutions in neutral state-dependent delays equations and models, *J. Comput. Appl. Math.* 174 (1) (2005) 179–199.
- [14] E. Hernández, A. Prokopczyk, Luiz Ladeira, A note on partial functional differential equations with state-dependent delay, *Nonlinear Anal. RWA* 7 (4) (2006) 510–519.
- [15] A.V. Rezounenko, Jianhong Wu, A non-local PDE model for population dynamics with state-selective delay: local theory and global attractors, *J. Comput. Appl. Math.* 190 (1–2) (2006) 99–113.
- [16] J.K. Hale, Verduyn Lunel, *Introduction to functional–differential equations*, *Applied Mathematical Sciences*, vol. 99, Springer-Verlag, New York, 1993.
- [17] M. Adimy, K. Ezzinbi, A class of linear partial neutral functional–differential equations with nondense domain, *J. Differen. Equat.* 147 (2) (1998) 285–332.
- [18] J.K. Hale, Partial neutral functional–differential equations, *Rev. Roumaine Math. Pures Appl.* 39 (4) (1994) 339–344.
- [19] J. Wu, H. Xia, Self-sustained oscillations in a ring array of coupled lossless transmission lines, *J. Differen. Equat.* 124 (1) (1996) 247–278.
- [20] Jianhong Wu, *Theory and applications of partial functional–differential equations*, *Applied Mathematical Sciences*, vol. 119, Springer-Verlag, New York, 1996.
- [21] E. Hernández, H. Henríquez, Existence results for partial neutral functional differential equations with unbounded delay, *J. Math. Anal. Appl.* 221 (2) (1998) 452–475.
- [22] E. Hernández, H. Henríquez, Existence of periodic solutions of partial neutral functional differential equations with unbounded delay, *J. Math. Anal. Appl.* 221 (2) (1998) 499–522.
- [23] E. Hernández, Existence results for partial neutral integrodifferential equations with unbounded delay, *J. Math. Anal. Appl.* 292 (1) (2004) 194–210.
- [24] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, *Applied Mathematical Sciences*, vol. 44, Springer-Verlag, New York, 1983.
- [25] Yoshiyuki Hino, Satoru Murakami, Toshiki Naito, *Functional–differential equations with infinite delay*, *Lecture Notes in Mathematics*, vol. 1473, Springer-Verlag, Berlin, 1991.
- [26] A. Granas, J. Dugundji, *Fixed Point Theory*, Springer-Verlag, New York, 2003.
- [27] R.H. Martin, *Nonlinear Operators and Differential Equations in Banach Spaces*, Robert E. Krieger Publ. Co., Florida, 1987.