

West Chester University

## Digital Commons @ West Chester University

---

Mathematics Faculty Publications

Mathematics

---

2008

### Some identities between basic hypergeometric series deriving from a new Bailey-type transformation

James McLaughlin

West Chester University of Pennsylvania, jmclaughlin2@wcupa.edu

Peter Zimmer

West Chester University of Pennsylvania, pzimmer@wcupa.edu

Follow this and additional works at: [https://digitalcommons.wcupa.edu/math\\_facpub](https://digitalcommons.wcupa.edu/math_facpub)



Part of the [Number Theory Commons](#)

---

#### Recommended Citation

McLaughlin, J., & Zimmer, P. (2008). Some identities between basic hypergeometric series deriving from a new Bailey-type transformation. *Journal of Mathematical Analysis and Applications*, 345(2), 670-677. Retrieved from [https://digitalcommons.wcupa.edu/math\\_facpub/53](https://digitalcommons.wcupa.edu/math_facpub/53)

This Article is brought to you for free and open access by the Mathematics at Digital Commons @ West Chester University. It has been accepted for inclusion in Mathematics Faculty Publications by an authorized administrator of Digital Commons @ West Chester University. For more information, please contact [wccressler@wcupa.edu](mailto:wccressler@wcupa.edu).

# SOME IDENTITIES BETWEEN BASIC HYPERGEOMETRIC SERIES DERIVING FROM A NEW BAILEY-TYPE TRANSFORMATION

JAMES MC LAUGHLIN AND PETER ZIMMER

ABSTRACT. We prove a new Bailey-type transformation relating WP-Bailey pairs. We then use this transformation to derive a number of new 3- and 4-term transformation formulae between basic hypergeometric series.

## 1. INTRODUCTION

Bailey's transform can be stated as follows:

**Lemma 1.** *Subject to suitable convergence conditions, if*

$$\beta_n = \sum_{r=0}^n \alpha_r U_{n-r} V_{n+r} \quad \text{and} \quad \gamma_n = \sum_{r=n}^{\infty} \delta_r U_{r-n} V_{r+n},$$

then

$$\sum_{n=0}^{\infty} \alpha_n \gamma_n = \sum_{n=0}^{\infty} \beta_n \delta_n.$$

The proof follows by switching the order of summation (see [2], pages 583–584, for example). Bailey set

$$U_n = \frac{1}{(q; q)_n}, \quad V_n = \frac{1}{(x; q)_n}, \quad \delta_n = (y, z; q)_n \left( \frac{x}{yz} \right)^n,$$

and used the  $q$ -Gauss sum,

$$(1.1) \quad {}_2\phi_1(a, b; c; q, c/ab) = \frac{(c/a, c/b; q)_{\infty}}{(c, c/ab; q)_{\infty}},$$

to get that

$$(1.2) \quad \sum_{n=0}^{\infty} (y, z; q)_n \left( \frac{x}{yz} \right)^n \beta_n = \frac{(x/y, x/z; q)_{\infty}}{(x, x/yz; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(y, z; q)_n}{(x/y, x/z; q)_n} \left( \frac{x}{yz} \right)^n \alpha_n,$$

---

*Date:* April 17, 2008.

*2000 Mathematics Subject Classification.* Primary: 33D15. Secondary: 11B65, 05A19.

*Key words and phrases.* Q-Series, basic hypergeometric series, Bailey chains, Bailey transform, WP-Bailey pairs .

where  $\alpha_0 = 1$  and

$$(1.3) \quad \beta_n = \sum_{r=0}^n \frac{\alpha_r}{(q; q)_{n-r} (x; q)_{n+r}}.$$

Here we are employing the usual notations. Let  $a$  and  $q$  be complex numbers, with  $|q| < 1$  unless otherwise stated. Then

$$(a)_0 = (a; q)_0 := 1, \quad (a)_n = (a; q)_n := \prod_{j=0}^{n-1} (1 - aq^j), \text{ for } n \in \mathbb{N},$$

$$(a_1; q)_n (a_2; q)_n \cdots (a_k; q)_n = (a_1, a_2, \dots, a_k; q)_n,$$

$$(a; q)_\infty := \prod_{j=0}^{\infty} (1 - aq^j),$$

$$(a_1; q)_\infty (a_2; q)_\infty \cdots (a_k; q)_\infty = (a_1, a_2, \dots, a_k; q)_\infty.$$

An  ${}_r\phi_s$  basic hypergeometric series is defined by

$${}_r\phi_s \left[ \begin{matrix} a_1, a_2, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, x \right] = \sum_{n=0}^{\infty} \frac{(a_1; q)_n (a_2; q)_n \cdots (a_r; q)_n}{(q; q)_n (b_1; q)_n \cdots (b_s; q)_n} \left( (-1)^n q^{n(n-1)/2} \right)^{s+1-r} x^n.$$

In modern notation, the pair of sequences  $(\alpha_n, \beta_n)$  above are termed a *Bailey pair relative to  $x/q$* . Slater, in [10] and [11], subsequently used this transformation of Bailey to derive 130 identities of the Rogers-Ramanujan type.

The first major variations in Bailey's construct at (1.2) appear to be due to Bressoud [5]. Another variation was given by Singh in [9]. All of these variations were put in a more formal setting by Andrews in [1], where he introduced a generalization of the standard Bailey pair as defined at (1.3) (see also equation (9.3) in [4]).

**Definition.** (Andrews [1]) Two sequences  $(\alpha_n(a, k), \beta_n(a, k))$  form a *WP-Bailey pair* provided

$$(1.4) \quad \beta_n(a, k) = \sum_{j=0}^n \frac{(k/a)_{n-j} (k)_{n+j}}{(q)_{n-j} (aq)_{n+j}} \alpha_j(a, k).$$

Note that if  $k = 0$ , then the definition reverts to that of a standard Bailey pair.

In the same paper Andrews showed that there were two distinct ways to construct new WP-Bailey pairs from a given pair. If  $(\alpha_n(a, k), \beta_n(a, k))$  satisfy (1.4), then so do  $(\alpha'_n(a, k), \beta'_n(a, k))$  and  $(\tilde{\alpha}_n(a, k), \tilde{\beta}_n(a, k))$ , where

$$(1.5) \quad \alpha'_n(a, k) = \frac{(\rho_1, \rho_2)_n}{(aq/\rho_1, aq/\rho_2)_n} \left( \frac{k}{c} \right)^n \alpha_n(a, c),$$

$$\beta'_n(a, k) = \frac{(k\rho_1/a, k\rho_2/a)_n}{(aq/\rho_1, aq/\rho_2)_n} \times \sum_{j=0}^n \frac{(1 - cq^{2j})(\rho_1, \rho_2)_j (k/c)_{n-j} (k)_{n+j}}{(1 - c)(k\rho_1/a, k\rho_2/a)_n (q)_{n-j} (qc)_{n+j}} \left(\frac{k}{c}\right)^j \beta_j(a, c),$$

with  $c = k\rho_1\rho_2/aq$  for the pair above, and

$$(1.6) \quad \begin{aligned} \tilde{\alpha}_n(a, k) &= \frac{(qa^2/k)_{2n}}{(k)_{2n}} \left(\frac{k^2}{qa^2}\right)^n \alpha_n\left(a, \frac{qa^2}{k}\right), \\ \tilde{\beta}_n(a, k) &= \sum_{j=0}^n \frac{(k^2/qa^2)_{n-j}}{(q)_{n-j}} \left(\frac{k^2}{qa^2}\right)^j \beta_j\left(a, \frac{qa^2}{k}\right). \end{aligned}$$

Andrews two constructions can be shown to imply the following Bailey-type transformations for WP-Bailey pairs, assuming suitable convergence conditions.

**Theorem 1.** *If  $(\alpha_n(a, k), \beta_n(a, k))$  satisfy*

$$\beta_n(a, k) = \sum_{j=0}^n \frac{(k/a)_{n-j} (k)_{n+j}}{(q)_{n-j} (aq)_{n+j}} \alpha_j(a, k),$$

then

$$(1.7) \quad \begin{aligned} \sum_{n=0}^{\infty} \frac{(1 - kq^{2n})(\rho_1, \rho_2; q)_n}{(1 - k)(kq/\rho_1, kq/\rho_2; q)_n} \left(\frac{aq}{\rho_1\rho_2}\right)^n \beta_n(a, k) = \\ \frac{(kq, kq/\rho_1\rho_2, aq/\rho_1, aq/\rho_2; q)_{\infty}}{(kq/\rho_1, kq/\rho_2, aq/\rho_1\rho_2, aq; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(\rho_1, \rho_2; q)_n}{(aq/\rho_1, aq/\rho_2; q)_n} \left(\frac{aq}{\rho_1\rho_2}\right)^n \alpha_n(a, k), \end{aligned}$$

and

$$(1.8) \quad \begin{aligned} \sum_{n=0}^{\infty} \left(\frac{qa^2}{k^2}\right)^n \beta_n(a, k) \\ = \frac{(qa/k, qa^2/k; q)_{\infty}}{(qa, qa^2/k^2; q)_{\infty}} \sum_{n=0}^{\infty} \frac{(k; q)_{2n}}{(qa^2/k; q)_{2n}} \left(\frac{qa^2}{k^2}\right)^n \alpha_n(a, k). \end{aligned}$$

We had initially derived the transformation at (1.7) in a way that was similar to the way Bailey derived (1.2), before finding that it followed from Andrews' first construction at (1.5). A result equivalent to the transformation at (1.8) was also stated by Bressoud in [5].

In the present paper we prove the following transformation for WP-Bailey pairs.

**Theorem 2.** *Subject to suitable convergence conditions, if*

$$(1.9) \quad \beta_n = \sum_{r=0}^n \frac{(k/a; q)_{n-r}}{(q; q)_{n-r}} \frac{(k; q)_{n+r}}{(aq; q)_{n+r}} \alpha_r,$$

then

$$\begin{aligned}
(1.10) \quad & \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty} \\
& \times \sum_{n=0}^{\infty} \frac{(q\sqrt{k}, -q\sqrt{k}, k^2/ab, b, \sqrt{qa}, -\sqrt{qa}; q)_n}{(\sqrt{k}, -\sqrt{k}, qab/k, kq/b, k\sqrt{q/a}, -k\sqrt{q/a}; q)_n} \left(\frac{-qa}{k}\right)^n \beta_n \\
& = \left(\frac{qk^2}{ab}, bq, \frac{q^2a^2b}{k^2}, \frac{q^2a}{b}; q^2\right)_\infty \sum_{n=0}^{\infty} \frac{\left(\frac{k^2}{ab}, b; q^2\right)_n}{\left(\frac{q^2a^2b}{k^2}, \frac{q^2a}{b}; q^2\right)_n} \left(\frac{-qa}{k}\right)^{2n} \alpha_{2n} \\
& \quad + \left(\frac{k^2}{ab}, b, \frac{q^3a^2b}{k^2}, \frac{q^3a}{b}; q^2\right)_\infty \sum_{n=0}^{\infty} \frac{\left(\frac{k^2q}{ab}, bq; q^2\right)_n}{\left(\frac{q^3a^2b}{k^2}, \frac{q^3a}{b}; q^2\right)_n} \left(\frac{-qa}{k}\right)^{2n+1} \alpha_{2n+1}.
\end{aligned}$$

We use this transformation to derive some new 3- and 4-term transformations between basic hypergeometric series.

## 2. A TRANSFORMATION DERIVING FROM A $q$ -ANALOG OF WATSON'S ${}_3F_2$ SUM

We recall the following  $q$ -analogue of Watson's  ${}_3F_2$  sum (see [6, (II.16), page 355]),

$$\begin{aligned}
(2.1) \quad & {}_8\phi_7 \left[ \lambda, q\sqrt{\lambda}, -q\sqrt{\lambda}, a, b, \lambda\sqrt{q/ab}, -\lambda\sqrt{q/ab}, ab/\lambda; q, -\frac{q\lambda}{ab} \right] \\
& = \frac{(\lambda q, \lambda q/ab; q)_\infty (aq, bq, q^2\lambda^2/a^2b, q^2\lambda^2/ab^2; q^2)_\infty}{(\lambda q/a, \lambda q/b; q)_\infty (q, abq, q^2\lambda^2/ab, q^2\lambda^2/a^2b^2; q^2)_\infty}.
\end{aligned}$$

We now prove the main theorem.

*Proof of Theorem 2.* In Lemma 1, set

$$\begin{aligned}
U_r &= \frac{(ab/\lambda; q)_r}{(q; q)_r}, \\
V_r &= \frac{(\lambda; q)_r}{(\lambda^2q/ab; q)_r}, \\
\delta_r &= \frac{(q\sqrt{\lambda}, -q\sqrt{\lambda}, a, b, \lambda\sqrt{q/ab}, -\lambda\sqrt{q/ab}; q)_r}{(\sqrt{\lambda}, -\sqrt{\lambda}, \lambda q/a, \lambda q/b, \sqrt{qab}, -\sqrt{qab}; q)_r} \left(\frac{-q\lambda}{ab}\right)^r.
\end{aligned}$$

Then

$$\begin{aligned}
\gamma_n &= \sum_{r=n}^{\infty} \delta_r U_{r-n} V_{r+n} \\
&= \sum_{m=0}^{\infty} \delta_{m+n} U_m V_{m+2n}
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{m=0}^{\infty} \frac{(q\sqrt{\lambda}, -q\sqrt{\lambda}, a, b, \lambda\sqrt{q/ab}, -\lambda\sqrt{q/ab}; q)_{m+n} (ab/\lambda; q)_m}{(\sqrt{\lambda}, -\sqrt{\lambda}, \lambda q/a, \lambda q/b, \sqrt{qab}, -\sqrt{qab}; q)_{m+n} (q; q)_m} \\
 &\quad \times \frac{(\lambda; q)_{m+2n}}{(\lambda^2 q/ab; q)_{m+2n}} \left( \frac{-q\lambda}{ab} \right)^{m+n} \\
 &= \frac{(q\sqrt{\lambda}, -q\sqrt{\lambda}, a, b, \lambda\sqrt{q/ab}, -\lambda\sqrt{q/ab}; q)_n (\lambda; q)_{2n}}{(\sqrt{\lambda}, -\sqrt{\lambda}, \lambda q/a, \lambda q/b, \sqrt{qab}, -\sqrt{qab}; q)_n (\lambda^2 q/ab; q)_{2n}} \left( \frac{-q\lambda}{ab} \right)^n \\
 &\times \sum_{m=0}^{\infty} \frac{(q^{1+n}\sqrt{\lambda}, -q^{1+n}\sqrt{\lambda}, aq^n, bq^n, \lambda\sqrt{q/ab}q^n, -\lambda\sqrt{q/ab}q^n; q)_m}{(\sqrt{\lambda}q^n, -\sqrt{\lambda}q^n, \lambda q^{1+n}/a, \lambda q^{1+n}/b, \sqrt{qab}q^n, -\sqrt{qab}q^n; q)_m} \\
 &\quad \times \frac{(\lambda q^{2n}, ab/\lambda; q)_m}{(\lambda^2 q^{1+2n}/ab, q; q)_m} \left( \frac{-q\lambda}{ab} \right)^m \\
 &= \frac{(q\sqrt{\lambda}, -q\sqrt{\lambda}, a, b, \lambda\sqrt{q/ab}, -\lambda\sqrt{q/ab}; q)_n (\lambda; q)_{2n}}{(\sqrt{\lambda}, -\sqrt{\lambda}, \lambda q/a, \lambda q/b, \sqrt{qab}, -\sqrt{qab}; q)_n (\lambda^2 q/ab; q)_{2n}} \left( \frac{-q\lambda}{ab} \right)^n \\
 &\times \frac{(\lambda q^{1+2n}, \lambda q/ab; q)_{\infty}}{(\lambda q^{1+n}/a, \lambda q^{1+n}/b; q)_{\infty}} \frac{(aq^{1+n}, bq^{1+n}, q^{2+n}\lambda^2/a^2b, q^{2+n}\lambda^2/ab^2; q^2)_{\infty}}{(q, abq^{1+2n}, q^2\lambda^{2+2n}/ab, q^2\lambda^2/a^2b^2; q^2)_{\infty}} \\
 &= \frac{(\lambda q, \lambda q/ab; q)_{\infty}}{(\lambda q/a, \lambda q/b, q)_{\infty}} \frac{(aq^{1+n}, bq^{1+n}, \lambda^2 q^{2+n}/a^2b, \lambda^2 q^{2+n}/ab^2; q^2)_{\infty}}{(q, qab, \lambda^2 q^2/ab, \lambda^2 q^2/a^2b^2; q^2)_{\infty}} \\
 &\quad \times (a, b; q)_n \left( \frac{-q\lambda}{ab} \right)^n \\
 &= \frac{(\lambda q, \lambda q/ab; q)_{\infty}}{(\lambda q/a, \lambda q/b, q)_{\infty}} \left( \frac{-q\lambda}{ab} \right)^n \times \\
 &\begin{cases} \frac{(aq, bq, \lambda^2 q^2/a^2b, \lambda^2 q^2/ab^2; q^2)_{\infty}}{(q, qab, \lambda^2 q^2/ab, \lambda^2 q^2/a^2b^2; q^2)_{\infty}} \frac{(a, b; q^2)_{\frac{n}{2}}}{(\lambda^2 q^2/a^2b, \lambda^2 q^2/ab^2; q^2)_{\frac{n}{2}}}, & n \text{ even,} \\ \frac{(a, b, \lambda^2 q^3/a^2b, \lambda^2 q^3/ab^2; q^2)_{\infty}}{(q, qab, \lambda^2 q^2/ab, \lambda^2 q^2/a^2b^2; q^2)_{\infty}} \frac{(aq, bq; q^2)_{\frac{n-1}{2}}}{(\lambda^2 q^3/a^2b, \lambda^2 q^3/ab^2; q^2)_{\frac{n-1}{2}}}, & n \text{ odd.} \end{cases}
 \end{aligned}$$

The fifth equality comes from applying (2.1) to the sum from the line before, after replacing  $\lambda$  with  $\lambda q^{2n}$ ,  $a$  with  $aq^n$  and  $b$  with  $bq^n$  in this identity.

We next make the substitutions  $\lambda \rightarrow k$ ,  $a \rightarrow k^2/bc$  followed by  $c \rightarrow a$ , and again suppose the sequences  $\{\alpha_n\}$  and  $\{\beta_n\}$  are related by

$$\beta_n = \sum_{r=0}^n \alpha_r U_{n-r} V_{n+r} = \sum_{r=0}^n \frac{(k/a; q)_{n-r}}{(q; q)_{n-r}} \frac{(k; q)_{n+r}}{(aq; q)_{n+r}} \alpha_r.$$

The result now follows.  $\square$

We can use this theorem to re-derive some known identities between basic hypergeometric series, and also to derive some new identities. For example,

inserting the “trivial” WP-Bailey pair

$$\alpha_n(a, k) = \begin{cases} 1 & n = 0, \\ 0, & n > 0, \end{cases}$$

$$\beta_n(a, k) = \frac{(k/a, k; q)_n}{(q, aq; q)_n}$$

in (1.10) leads to a version of (2.1) (perhaps not surprisingly, since (2.1) was used to prove Theorem 2). We also note in passing that applying Andrews first construction at (1.5) to this trivial WP-Bailey pair leads to a variant of Jackson’s sum of a terminating  ${}_8\phi_7$ , while applying his second construction at (1.6) leads to a variant of the  $q$ -Pfaff-Saalschütz sum.

Before going further, we introduce some standard space-saving notation:

$${}_{r+1}W_r(a_1; a_4, \dots, a_{r+1}; q, z) = {}_{r+1}\phi_r \left[ \begin{matrix} a_1, q\sqrt{a_1}, -q\sqrt{a_1}, a_4, \dots, a_{r+1} \\ \sqrt{a_1}, -\sqrt{a_1}, \frac{a_1q}{a_4}, \dots, \frac{a_1q}{a_{r+1}} \end{matrix}; q, z \right].$$

Inserting the “unit” WP-Bailey pair (see [3] for example, where this WP-Bailey pair, and others employed below, may be found),

$$\alpha_n(a, k) = \frac{(q\sqrt{a}, -q\sqrt{a}, a, a/k; q)_n}{(\sqrt{a}, -\sqrt{a}, q, kq; q)_n} \left(\frac{k}{a}\right)^n,$$

$$\beta_n(a, k) = \begin{cases} 1 & n = 0, \\ 0, & n > 1, \end{cases}$$

in (1.10) and replacing  $q$  with  $\sqrt{q}$  leads to the following identity:

$$\frac{\left(\frac{\sqrt{q}ab}{k}, \frac{qab}{k}, \frac{k\sqrt{q}}{b}, \frac{kq}{b}, -\frac{a\sqrt{q}}{k}, -\frac{aq}{k}, \sqrt{q}, \frac{k^2\sqrt{q}}{a}, qa; q\right)_{\infty}}{(kq, k\sqrt{q}; q)_{\infty}}$$

$$= \left(\frac{k^2\sqrt{q}}{ab}, b\sqrt{q}, \frac{a^2bq}{k^2}, \frac{qa}{b}; q\right)_{\infty} {}_8W_7 \left(a; a\sqrt{q}, \frac{a}{k}, \frac{a\sqrt{q}}{k}, \frac{k^2}{ab}, b; q, q\right)$$

$$- q^{1/2} \frac{(1-aq)(1-a/k)}{(1-\sqrt{q})(1-k\sqrt{q})} \left(\frac{k^2}{ab}, b, \frac{a^2bq^{3/2}}{k^2}, \frac{q^{3/2}a}{b}; q\right)_{\infty}$$

$$\times {}_8W_7 \left(aq; a\sqrt{q}, \frac{a\sqrt{q}}{k}, \frac{aq}{k}, \frac{k^2\sqrt{q}}{ab}, b\sqrt{q}; q, q\right).$$

This identity is a particular case of Bailey’s nonterminating extension of Jackson’s  ${}_8\phi_7$  sum (see [6, page 356, II.25]).

We now state some transformations which we believe are new. Upon substituting Singh’s WP-Bailey pair [9] (see also [3], where one of Andrews’

constructions was used to derive Singh's pair from the unit pair),

$$(2.2) \quad \alpha_n(a, k) = \frac{(q\sqrt{a}, -q\sqrt{a}, a, y, z, a^2q/kyz; q)_n}{(\sqrt{a}, -\sqrt{a}, q, aq/y, aq/z, kyz/a; q)_n} \left(\frac{k}{a}\right)^n,$$

$$\beta_n(a, k) = \frac{(ky/a, kz/a, k, aq/yz; q)_n}{(aq/y, aq/z, kyz/a, q; q)_n},$$

into (1.10), we get the following transformation.

**Corollary 1.**

$$(2.3) \quad \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty}$$

$$\times {}_{10}W_9 \left( k; \frac{k^2}{ab}, b, \sqrt{qa}, -\sqrt{qa}, \frac{ky}{a}, \frac{kz}{a}, \frac{aq}{yz}; q, -\frac{qa}{k} \right)$$

$$= \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty$$

$$\times {}_{12}W_{11} \left( a; \frac{k^2}{ab}, b, aq, y, yq, z, zq, \frac{a^2q}{kyz}, \frac{a^2q^2}{kyz}; q^2, q^2 \right)$$

$$- q \frac{(1-aq^2)(1-y)(1-z)(1-a^2q/kyz)}{(1-q)(1-aq/y)(1-aq/z)(1-kyz/a)} \left( \frac{k^2}{ab}, b, \frac{a^2bq^3}{k^2}, \frac{q^3a}{b}; q^2 \right)_\infty$$

$$\times {}_{12}W_{11} \left( q^2a; \frac{k^2q}{ab}, bq, aq, yq, yq^2, zq, zq^2, \frac{a^2q^2}{kyz}, \frac{a^2q^3}{kyz}; q^2, q^2 \right).$$

Remark: The identity above may be regarded as an extension of (2.1), since substituting  $y = 1$  in (2.3) leads to a variant of (2.1).

We next apply the theorem to some WP-Bailey pairs found by Andrews and Berkovich [3].

**Corollary 2.**

$$(2.4) \quad \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty}$$

$$\times {}_7\phi_6 \left[ \begin{matrix} q\sqrt{k}, -q\sqrt{k}, \frac{k^2}{ab}, b, \sqrt{qa}, -\sqrt{qa}, \frac{k^2}{qa^2} \\ \sqrt{k}, -\sqrt{k}, \frac{qab}{k}, \frac{qk}{b}, k\sqrt{\frac{q}{a}}, -k\sqrt{\frac{q}{a}} \end{matrix}; q, -\frac{qa}{k} \right]$$

$$= \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty$$

$$\times {}_{16}W_{15} \left( a; \frac{k^2}{ab}, b, aq, \frac{k}{aq}, \frac{k}{a}, a\sqrt{\frac{q}{k}}, -a\sqrt{\frac{q}{k}}, \frac{aq}{\sqrt{k}}, -\frac{aq}{\sqrt{k}}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}, \frac{aq^2}{\sqrt{k}}, -\frac{aq^2}{\sqrt{k}}; q^2, q^2 \right)$$

$$- q \frac{(1-aq^2)(1-qa^2/k)(1-q^2a^2/k)(1-k/aq)}{(1-q)(1-k)(1-kq)(1-a^2q^2/k)} \left( \frac{k^2}{ab}, b, \frac{a^2bq^3}{k^2}, \frac{q^3a}{b}; q^2 \right)_\infty \times$$



$${}_{16}W_{15} \left( aq^2; \frac{k^2q}{ab}, bq, aq, \frac{kq}{a}, \frac{k}{a}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}, \frac{aq^2}{\sqrt{k}}, \frac{-aq^2}{\sqrt{k}}, a\sqrt{\frac{q^5}{k}}, -a\sqrt{\frac{q^5}{k}}, \frac{aq^3}{\sqrt{k}}, \frac{-aq^3}{\sqrt{k}}; q^2, q^2 \right).$$

*Proof.* Insert the WP-Bailey pair

$$\begin{aligned} \alpha_n(a, k) &= \frac{(a, q\sqrt{a}, -q\sqrt{a}, k/aq; q)_n (qa^2/k; q)_{2n}}{(q, \sqrt{a}, -\sqrt{a}, a^2q^2/k; q)_n (k; q)_{2n}} \left( \frac{k}{a} \right)^n, \\ \beta_n(a, k) &= \frac{(k^2/qa^2; q)_n}{(q; q)_n}, \end{aligned}$$

from [3] into (1.10).  $\square$

**Corollary 3.**

$$\begin{aligned} (2.5) \quad & \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty} \\ & \times {}_6\phi_5 \left[ \begin{matrix} -q\sqrt{k}, \frac{k^2}{ab}, b, \sqrt{qa}, -\sqrt{qa}, \frac{k^2}{a^2} \\ -\sqrt{k}, \frac{qab}{k}, \frac{qk}{b}, k\sqrt{\frac{q}{a}}, -k\sqrt{\frac{q}{a}} \end{matrix}; q, -\frac{qa}{k} \right] \\ & = \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty \times \\ & {}_{16}W_{15} \left( a; \frac{k^2}{ab}, b, aq, \frac{kq}{a}, \frac{k}{a}, a\sqrt{\frac{q}{k}}, -a\sqrt{\frac{q}{k}}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}, \frac{a}{\sqrt{k}}, \frac{aq}{\sqrt{k}}, -\frac{aq}{\sqrt{k}}, -\frac{aq^2}{\sqrt{k}}; q^2, q^2 \right) \\ & - q \frac{(1-aq^2)(1-\frac{k}{a})(1-\frac{a}{\sqrt{k}})(1+\frac{aq}{\sqrt{k}})}{(1-q)(1-kq)(1-\sqrt{k}q)(1+\sqrt{k})} \left( \frac{k^2}{ab}, b, \frac{a^2bq^3}{k^2}, \frac{q^3a}{b}; q^2 \right)_\infty \times \\ & {}_{16}W_{15} \left( aq^2; \frac{k^2q}{ab}, bq, aq, \frac{kq}{a}, \frac{kq^2}{a}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}, a\sqrt{\frac{q^5}{k}}, -a\sqrt{\frac{q^5}{k}}, \frac{aq}{\sqrt{k}}, \frac{aq^2}{\sqrt{k}}, \frac{-aq^2}{\sqrt{k}}, \frac{-aq^3}{\sqrt{k}}; q^2, q^2 \right). \end{aligned}$$

*Proof.* This transformation follows similarly, after inserting the WP-Bailey pair

$$\begin{aligned} \alpha_n(a, k) &= \frac{(a, q\sqrt{a}, -q\sqrt{a}, a\sqrt{\frac{q}{k}}, -a\sqrt{\frac{q}{k}}, \frac{a}{\sqrt{k}}, -\frac{aq}{\sqrt{k}}, \frac{k}{a}; q)_n}{(q, \sqrt{a}, -\sqrt{a}, \sqrt{qk}, -\sqrt{qk}, q\sqrt{k}, -\sqrt{k}, \frac{qa^2}{k}; q)_n} \left( \frac{k}{a} \right)^n, \\ \beta_n(a, k) &= \frac{(\sqrt{k}, \frac{k^2}{a^2}; q)_n}{(q, q\sqrt{k}; q)_n}, \end{aligned}$$

from [3] into (1.10).  $\square$

We next consider two WP-Bailey pairs found by Bressoud [5] (see also [3], where these pairs are also investigated). We do not consider Bressoud's first WP-Bailey pair, since, as remarked in [3], it is a limiting case of Singh's WP-Bailey pair at (2.2).

**Corollary 4.**

$$\begin{aligned}
(2.6) \quad & \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty} \\
& \times {}_8W_7 \left( k; \frac{k^2}{ab}, b, \sqrt{aq}, \frac{aq}{k}, \frac{-k}{\sqrt{a}}; q, -\sqrt{q} \right) \\
& = \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty \\
& \quad \times {}_{10}W_9 \left( \sqrt{a}; \frac{k}{\sqrt{ab}}, \frac{-k}{\sqrt{ab}}, \sqrt{b}, -\sqrt{b}, \sqrt{aq}, \frac{a\sqrt{q}}{k}, \frac{aq}{k}; q, q \right) \\
& \quad - \sqrt{q} \frac{(1 - q\sqrt{a}) \left(1 - \frac{a\sqrt{q}}{k}\right)}{(1 - \sqrt{q})(1 - \frac{k}{\sqrt{a}})} \left( \frac{k^2}{ab}, b, \frac{a^2bq^3}{k^2}, \frac{q^3a}{b}; q^2 \right)_\infty \\
& \quad \times {}_{10}W_9 \left( q\sqrt{a}; k\sqrt{\frac{q}{ab}}, -k\sqrt{\frac{q}{ab}}, \sqrt{bq}, -\sqrt{bq}, \sqrt{aq}, \frac{a\sqrt{q^3}}{k}, \frac{aq}{k}; q, q \right).
\end{aligned}$$

*Proof.* Insert Bressoud's second WP-Bailey pair

$$\begin{aligned}
\alpha_n(a, k) &= \frac{1 - \sqrt{a}q^n}{1 - \sqrt{a}} \frac{\left(\sqrt{a}, \frac{a\sqrt{q}}{k}; \sqrt{q}\right)_n}{\left(\sqrt{q}, \frac{k}{\sqrt{a}}; \sqrt{q}\right)_n} \left(\frac{k}{a\sqrt{q}}\right)^n, \\
\beta_n(a, k) &= \frac{\left(k, \frac{aq}{k}; q\right)_n}{\left(q, \frac{k^2}{a}; q\right)_n} \frac{\left(\frac{-k}{\sqrt{a}}; \sqrt{q}\right)_{2n}}{\left(-\sqrt{aq}; \sqrt{q}\right)_{2n}} \left(\frac{k}{a\sqrt{q}}\right)^n,
\end{aligned}$$

into (1.10). □

**Corollary 5.**

$$\begin{aligned}
(2.7) \quad & \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty} \\
& \times {}_8W_7 \left( k; \frac{k^2}{ab}, b, \sqrt{aq}, \frac{a}{k}, \frac{-kq}{\sqrt{a}}; q, -\sqrt{q} \right) \\
& = \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty \times \\
& \quad {}_{12}W_{11} \left( \sqrt{a}; iq a^{1/4}, -iq a^{1/4}, \frac{k}{\sqrt{ab}}, \frac{-k}{\sqrt{ab}}, \sqrt{b}, -\sqrt{b}, \sqrt{aq}, \frac{a}{k}, \frac{a\sqrt{q}}{k}; q, q \right) \\
& \quad - \sqrt{q} \frac{(1 - aq^2) \left(1 - \frac{a}{k}\right)}{(1 - \sqrt{q}) \left(1 - k\sqrt{\frac{q}{a}}\right) (1 + \sqrt{a})} \left( \frac{k^2}{ab}, b, \frac{a^2bq^3}{k^2}, \frac{q^3a}{b}; q^2 \right)_\infty \times \\
& \quad {}_{12}W_{11} \left( q\sqrt{a}; iq^{3/2}a^{1/4}, -iq^{3/2}a^{1/4}, k\sqrt{\frac{q}{ab}}, -k\sqrt{\frac{q}{ab}}, \sqrt{bq}, -\sqrt{bq}, \sqrt{aq}, \frac{a\sqrt{q}}{k}, \frac{aq}{k}; q, q \right).
\end{aligned}$$

*Proof.* Insert Bressoud's third WP-Bailey pair

$$\alpha_n(a, k) = \frac{1 - aq^{2n}}{1 - a} \frac{(\sqrt{a}, \frac{a}{k}; \sqrt{q})_n}{(\sqrt{q}, k\sqrt{\frac{q}{a}}; \sqrt{q})_n} \left(\frac{k}{a\sqrt{q}}\right)^n,$$

$$\beta_n(a, k) = \frac{\left(k, \frac{a}{k}, -k\sqrt{\frac{q}{a}}, -\frac{kq}{\sqrt{a}}; q\right)_n}{\left(q, \frac{qk^2}{a}, -\sqrt{a}, -\sqrt{aq}; q\right)_n} \left(\frac{k}{a\sqrt{q}}\right)^n,$$

into (1.10).  $\square$

Finally, we apply the theorem to three WP-Bailey pairs found by the present authors in [8]:

$$(2.8) \quad \alpha_n^{(1)}(a, k) = \frac{(qa^2/k^2; q)_n}{(q, q)_n} \left(\frac{k}{a}\right)^n,$$

$$\beta_n^{(1)}(a, k) = \frac{(qa/k, k; q)_n (k^2/a; q)_{2n}}{(k^2/a, q, q)_n (aq, q)_{2n}}.$$

$$(2.9) \quad \alpha_n^{(2)}(a, k) = \frac{(a, q\sqrt{a}, -q\sqrt{a}, k/a, a\sqrt{q/k}, -a\sqrt{q/k}; q)_n}{(\sqrt{a}, -\sqrt{a}, qa^2/k, \sqrt{qk}, -\sqrt{qk}, q; q)_n} (-1)^n,$$

$$\beta_n^{(2)}(a, k) = \begin{cases} \frac{(k, k^2/a^2; q^2)_{n/2}}{(q^2, q^2a^2/k; q^2)_{n/2}}, & n \text{ even,} \\ 0, & n \text{ odd.} \end{cases}$$

$$(2.10) \quad \alpha_n^{(3)}(a, q) = \frac{(a, q\sqrt{a}, -q\sqrt{a}, d, q/d, -a; q)_n}{(\sqrt{a}, -\sqrt{a}, aq/d, ad, -q, q; q)_n} (-1)^n,$$

$$\beta_n^{(3)}(a, q) = \begin{cases} \frac{(q^2/ad, dq/a; q^2)_{n/2}}{(adq, aq^2/d; q^2)_{n/2}}, & n \text{ even,} \\ -a \frac{(q/ad, d/a; q^2)_{(n+1)/2}}{(ad, aq/d; q^2)_{(n+1)/2}}, & n \text{ odd.} \end{cases}$$

Note that the third pair is restricted in the sense that it is necessary to set  $k = q$  for (1.9) to hold.

**Corollary 6.**

$$(2.11) \quad \frac{(qab/k, kq/b; q)_\infty (q, k^2q/a, q^2a, q^2a^2/k^2; q^2)_\infty}{(kq, qa/k; q)_\infty}$$

$$\times {}_8W_7 \left( k; \frac{k^2}{ab}, b, \frac{qa}{k}, \frac{k}{\sqrt{a}}, \frac{-k}{\sqrt{a}}; q, \frac{-qa}{k} \right)$$

$$= \left( \frac{k^2q}{ab}, bq, \frac{a^2bq^2}{k^2}, \frac{q^2a}{b}; q^2 \right)_\infty {}_4\phi_3 \left[ \begin{matrix} \frac{k^2}{ab}, b, \frac{qa^2}{k^2}, \frac{q^2a^2}{k^2} \\ \frac{q^2a^2b}{k^2}, \frac{q^2a}{b}, q \end{matrix}; q^2, q^2 \right]$$

$$-q \frac{\left(1 - \frac{qa^2}{k^2}\right)}{(1-q)} \left(\frac{k^2}{ab}, b, \frac{a^2 bq^3}{k^2}, \frac{q^3 a}{b}; q^2\right)_{\infty} {}_4\phi_3 \left[ \begin{matrix} \frac{k^2 q}{ab}, bq, \frac{q^2 a^2}{k^2}, \frac{q^3 a^2}{k^2} \\ \frac{q^3 a^2 b}{k^2}, \frac{q^3 a}{b}, q^3 \end{matrix}; q^2, q^2 \right].$$

*Proof.* Insert the WP-Bailey pair at (2.8) into (1.10).  $\square$

Remark: The substitution  $k = a\sqrt{q}$  also gives a special case of (2.1).

**Corollary 7.**

$$(2.12) \quad \frac{(qab/k, kq/b; q)_{\infty} (q, k^2 q/a, q^2 a, q^2 a^2/k^2; q^2)_{\infty}}{(kq, qa/k; q)_{\infty}} \times$$

$${}_{12}W_{11} \left( k; \frac{k^2}{ab}, \frac{k^2 q}{ab}, b, bq, \sqrt{aq}, -\sqrt{aq}, \sqrt{aq^3}, -\sqrt{aq^3}, \frac{k^2}{a^2}; q^2, \frac{a^2 q^2}{k^2} \right)$$

$$= \left( \frac{k^2 q}{ab}, bq, \frac{a^2 bq^2}{k^2}, \frac{q^2 a}{b}; q^2 \right)_{\infty} \times$$

$${}_{12}W_{11} \left( a; \frac{k^2}{ab}, b, aq, \frac{k}{a}, \frac{kq}{a}, a\sqrt{\frac{q}{k}}, -a\sqrt{\frac{q}{k}}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}; q^2, \frac{q^2 a^2}{k^2} \right)$$

$$+ \frac{qa(1-aq^2)(1-\frac{k}{a})}{k(1-q)(1-kq)} \left( \frac{k^2}{ab}, b, \frac{a^2 bq^3}{k^2}, \frac{q^3 a}{b}; q^2 \right)_{\infty} \times$$

$${}_{12}W_{11} \left( aq^2; \frac{k^2 q}{ab}, bq, aq, \frac{kq}{a}, \frac{kq^2}{a}, a\sqrt{\frac{q^3}{k}}, -a\sqrt{\frac{q^3}{k}}, a\sqrt{\frac{q^5}{k}}, -a\sqrt{\frac{q^5}{k}}; q^2, \frac{q^2 a^2}{k^2} \right).$$

*Proof.* Insert the WP-Bailey pair at (2.9) into (1.10).  $\square$

**Corollary 8.**

$$(2.13) \quad \frac{\left(ab, \frac{q^2}{b}; q\right)_{\infty} \left(q, \frac{q^3}{a}, q^2 a, a^2; q^2\right)_{\infty}}{(q^2, a; q)_{\infty}} \times$$

$$\left[ {}_{14}W_{13} \left( q; \frac{q^2}{ab}, \frac{q^3}{ab}, b, bq, \sqrt{qa}, -\sqrt{qa}, \sqrt{q^3 a}, -\sqrt{q^3 a}, \frac{q^2}{ad}, \frac{qd}{a}, q^2; q^2, a^2 \right) \right.$$

$$+ a^2 \frac{(1-q^3) \left(1 - \frac{q^2}{ab}\right) (1-b)(1-aq) \left(1 - \frac{q}{ad}\right) \left(1 - \frac{d}{a}\right)}{(1-q) \left(1 - \frac{q^2}{b}\right) (1-ab)(1-ad) \left(1 - \frac{q^3}{a}\right) \left(1 - \frac{aq}{d}\right)} \times$$

$$\left. {}_{14}W_{13} \left( q^3; \frac{q^3}{ab}, \frac{q^4}{ab}, bq, bq^2, \sqrt{q^3 a}, -\sqrt{q^3 a}, \sqrt{q^5 a}, -\sqrt{q^5 a}, \frac{q^3}{ad}, \frac{q^2 d}{a}, q^2; q^2, a^2 \right) \right]$$

$$= \left( a^2 b, bq, \frac{q^3}{ab}, \frac{q^2 a}{b}; q^2 \right)_{\infty} {}_{12}W_{11} \left( a; \frac{q^2}{ab}, b, aq, -a, -aq, d, dq, \frac{q}{d}, \frac{q^2}{d}; q^2, a^2 \right)$$

$$+ a \frac{(1-aq^2)(1-d) \left(1 - \frac{q}{d}\right) (1+a)}{(1-q^2) \left(1 - \frac{aq}{d}\right) (1-ad)} \left( qa^2 b, b, \frac{q^2}{ab}, \frac{q^3 a}{b}; q^2 \right)_{\infty} \times$$

$${}_{12}W_{11} \left( aq^2; \frac{q^3}{ab}, bq, aq, -aq, -aq^2, dq, dq^2, \frac{q^2}{d}, \frac{q^3}{d}; q^2, a^2 \right).$$

*Proof.* Insert the WP-Bailey pair at (2.10) into (1.10), and set  $k = q$ .  $\square$

Remark: The extra  $q$ -products inserted in the numerators and denominators of the terms in the two series on the left in the identity above are there so as to give each these series the form of a  ${}_{r+1}W_r$  series.

#### REFERENCES

- [1] Andrews, George E. *Bailey's transform, lemma, chains and tree*. Special functions 2000: current perspective and future directions (Tempe, AZ), 1–22, NATO Sci. Ser. II Math. Phys. Chem., **30**, Kluwer Acad. Publ., Dordrecht, 2001.
- [2] Andrews, George E.; Askey, Richard; Roy, Ranjan *Special functions*. Encyclopedia of Mathematics and its Applications, 71. Cambridge University Press, Cambridge, 1999. xvi+664 pp.
- [3] Andrews, George; Berkovich, Alexander *The WP-Bailey tree and its implications*. J. London Math. Soc. (2) **66** (2002), no. 3, 529–549.
- [4] Bailey, W. N., *Some Identities in Combinatory Analysis*. Proc. London Math. Soc. **49** (1947) 421–435.
- [5] Bressoud, David, *Some identities for terminating  $q$ -series*. Math. Proc. Cambridge Philos. Soc. **89** (1981), no. 2, 211–223.
- [6] Gasper, George; Rahman, Mizan *Basic hypergeometric series*. With a foreword by Richard Askey. Second edition. Encyclopedia of Mathematics and its Applications, 96. Cambridge University Press, Cambridge, 2004. xxvi+428 pp.
- [7] Mc Laughlin, James; Zimmer, Peter. *Some Applications of a Bailey-type Transformation* - submitted
- [8] Mc Laughlin, James; Zimmer, Peter. *Some transformations for terminating basic hypergeometric series* - submitted
- [9] Singh, U. B. *A note on a transformation of Bailey*. Quart. J. Math. Oxford Ser. (2) **45** (1994), no. 177, 111–116.
- [10] Slater, L. J. *A new proof of Rogers's transformations of infinite series*. Proc. London Math. Soc. (2) **53**, (1951). 460–475.
- [11] Slater, L. J. *Further identities of the Rogers-Ramanujan type*, Proc. London Math. Soc. **54** (1952) 147–167.

MATHEMATICS DEPARTMENT, ANDERSON HALL, WEST CHESTER UNIVERSITY, WEST CHESTER, PA 19383

*E-mail address:* `jmclaughl@wcupa.edu`

MATHEMATICS DEPARTMENT, ANDERSON HALL, WEST CHESTER UNIVERSITY, WEST CHESTER, PA 19383

*E-mail address:* `pzimmer@wcupa.edu`